

maintain a steady level. On a seasonal basis, however, the monthly fluxes of most of the water quality variables exhibit significant mass during wet seasons and lower fluxes during dry seasons, except the ammonia flux, which exhibits an opposite trend. Further analyses demonstrate that the monthly mass fluxes are mainly controlled by the runoff; while the annual fluxes are mainly affected by the concentrations.

Rehabilitation and eco-environmental protection of the Yangtze Estuary

The Yangtze is the largest river in China, the world's famous "Treasure House of biota", and the "Golden Waterway" connecting the east, middle and west part of the nation. Located at the joint of the Yangtze Economic Zone and the Coastal Economic Zone, the Yangtze estuary occupies an essential position in China. Along with the fast economic growth in the past decades, the pollution discharge in the basin has been increasing rapidly leading the deterioration of water quality in the estuary region. As a result, the Yangtze estuary and its adjacent sea area has become one of the most serious places of "Red tide" occurring in west Pacific Ocean. After completion of the Three Gorges Project and the Water Transfer from South to North Project as well as a number of hydropower projects in the upper reaches, the temporal and spatial distribution of water resources in the basin will be alternated, which will exert significant impacts on the ecosystem and environment in the estuary region. In order to preserve the ecosystem and environment, the Rehabilitation Planning of the Yangtze Estuary has been compiled to protect the important wetland and bio-diversity in the region, control the water pollution and the salt-water intrusion to ensure the safety of water supply in the mean time of maintaining the deep channels for navigation.

Investigation of microbial diversity and its degradation potential to PAHs in the sediment in Bohai Bay, China, by PCR-DGGE

Liping WANG

River&Coastal Environment Research Center, Chinese Research Academy of Environmental Sciences, Beijing, 100012, China

In order to investigate the biodegradation potential of the indigenous microorganisms in the sediment in Bohai Bay, China, an improved understanding of the diversity and structure of the microbial communities in the sediment is required. Seventeen sediment samples were collected in Bohai Bay (117° 41' -117° 60' E, 38° 33' -39° 00' N). DGGE analysis of PCR-amplified 16S rDNA gene fragments confirmed that there was a remarkable different in the composition of the bacterial community in some different station. The

content of polycyclic aromatic hydrocarbons (PAHs) in the sediment was also investigated. The presence of common bands for microbial species in the native sediment DNA indicated that some strains could be potential in situ PAH-degraders.

Ecological characteristics of the macrobenthic community in the Changjiang Estuary and its adjacent waters

Lusan LIU

Chinese Research Academy of Environmental Science, Beijing 100012, China

E-mail: liuls@craes.org.cn

Spatial distribution and evolution of macrobenthos in the Changjiang estuary and its adjacent waters were presented in this study. Samples were collected from 86 stations in four investigations during 2005~2006. Among all the samples, a total of 330 macrobenthic species were identified including in the number of species: 122 Mollusca, 83 Polychaeta, 67 Crustacea, 23 Echinodermata, 28 Pisces and 7 other species. Total average abundance of macrobenthos in the study area was 146.4 ± 22.3 ind/m², the mean biomass (AFDW) was 2.31 ± 0.41 g·m⁻², and the mean secondary production was 2.48 ± 0.38 g (AFDW) m⁻² a⁻¹. The average values of Shannon-Wiener's, Margalef's and Pielou's indices were 1.72 ± 0.16 , 1.37 ± 0.19 , 0.64 ± 0.04 respectively.

The results showed that the macrobenthos of the study area could be divided into 3 levels from west to east, or from nearshore to offshore. In the most westward area named the Hekou subarea and the Hangzhou Bay subarea, the species composition of macrobenthos was very simple, and the indices of biodiversity were very low, indicating that the community structure was very susceptible. In the Kouwai subarea and the Zhoushan subarea close to the central part of the research region, the species composition of macrobenthos was more complex than that in the most westward area, and the indices of biodiversity were higher, indicating that the community structure was relatively steady. In the most outside area named the Jinhai subarea, the species composition was most complex in the whole research region, and the indices of biodiversity were highest, indicating that the community structure was most steady.

Comparisons of the present results with those of other sea areas in China make it clear that the macrobenthos usually has low production in estuary area where the environments are usually