

conducted, and potential cancer risk by PCBs and DDT was suggested. Based on the results, it was verified that POPs monitoring method with bivalves as a bioindicator is efficient and monitoring of PCBs and DDTs is significant in the Pearl River Delta.

Reference

Guo Jianyang, Eddy Y. Zeng¹, Wu Fengchang, et al (2007). Organochlorine Pesticides In Seafood Products From Southern China And Health Risk Assessment. *Env.Tox.Chem.*, 26: 1109-1115

Effect of the condensed tannin from the dead leaves of deciduous trees on the photosynthesis reaction inhibition to *Microcystis aeruginosa*

K SHIMADA^{1*}, N KITAMURA¹, S NOMOTO¹, Y YOSHIDA², H MATSUSHIMA² & YASADA³

¹ Department of Civil Engineering Graduate School of Science and Technology Nihon University

*E-Mail: yoshida@civil.cst.nihon-u.ac.jp

² Department of Civil Engineering College of Science and Technology Nihon University

³ Department of General Education College of Science and Technology Nihon University

Hepatotoxin like microcystin from *Microcystis aeruginosa* (*M.a.*) is extremely toxic to human and is an important water issue in terms of water resources and water quality preservations. The countermeasures to the water-bloom occurrence due to the excessive growth of toxic cyanobacteria are not necessarily available at present. The authors have revealed that the condensed tannin (CT) in the water-soluble extract from the dead leaves of deciduous trees is very effective to the growth control of *M.a.* Furthermore, the growth control was clearly affected by the condensed tannin loading intensity (CTLI). The mechanism of the *M.a.* growth control is unknown yet. Photosynthesis light reaction has been divided into NADP reduction reaction (PS I) and oxygen generation reaction (PS II). The addition of the extract results in the growth control and in fact decreases in the oxygen generation simultaneously, which imply the extract from the dead leaves functions like PS II inhibition. Accordingly, the effect of the photosynthesis reaction inhibition was examined on the basis of the direct measurement of the DO concentration trend in the *M.a.* culture solution to evaluate the oxygen generation rate (OGR). Data were plotted in Fig. 1 to show the photosynthesis inhibition effect on *M.a.* obtained from National Institute for Environmental Studies, Japan, (NIES) as NIES 102 when the CT was




Fig. 1 OGRs before(①) and after the extract addition(②)




Fig. 2 DO consumption by condensed tannin




Fig. 3 Effects of condensed tannin on RARP

added into the *M.a.* culture solution at about 2 mg/L CT concentration and at about 1.47×10^{-9} mg/cell CTLI. The *M.a.* culture solution containing about 0.9 million cells per ml was lighted so that the initial OGR for control was about 81.6 mg-O₂/L·h. The OGR in 2 minutes after the addition of the extract decreased immediately down to 27.6 mg-O₂/L·h. The extract itself was consuming the DO at about 0.72 mg-O₂/L·h, being very small and neglected when compared with the

actual OGR (Fig. 2). Consequently, the extract would be the major reason for the inhibition of the photosynthesis II reaction. In order to quantitatively evaluate the photosynthesis inhibition effect, the ratio of the OGRs before and after the extract addition was estimated as in terms of the residual activity ratio of the photosynthesis light reaction (RARP). Based on the data between RARP and CTLI, the RARP decreased clearly as the increase in the CTLI. A half-inhibition concentration (IC_{50}) is defined to be the critical CTLI value corresponding to the 50 % value of the RARP. Some IC_{50} values are given for *Prunus* as 1.9×10^{-9} mg/cell, for *Acer buergerianum* as 1.4×10^{-9} mg/cell, and for *Quercus acutissima* as 0.8×10^{-9} mg/cell in order, implying that the IC_{50} values would be depending on the sort of the deciduous trees or possibly on the extent of photosynthesis light reaction of the leaves of the trees.

The application of foraminifera and ostracod for long-term ecosystem monitoring in enclosed coastal sea: A case study of Osaka Bay, Seto Inland Sea, Japan

Akira TSUJIMOTO^{1*}, Moriaki YASUHARA²,
Shinichi NISHIDA¹, Hideo YAMAZAKI³,
Kotaro HIROSE¹ & Shusaku YOSHIKAWA¹

¹ Division of Biology and Geosciences, Graduate School of Science, Osaka City University, Osaka 558-8585, Japan

*E-mail: tujimoto@sci.osaka-cu.ac.jp

² Department of Paleobiology, National Museum of Natural History, Smithsonian Institution, Washington, DC 20013-7012, USA

³ Department of Life Science, School of Science and Engineering, Kinki University, Higashiosaka, Osaka 577-8502, Japan

Foraminifera and Ostracoda are shelled meiobenthos. They are sensitive to anthropogenic activities, and their remains are abundantly preserved as 'fossil' in bay sediments after their death. Thus, they can be used to trace environmental/ecosystem changes over periods of decades and centuries using sediment cores. In this study, we investigated spatial and temporal distributions of foraminiferal and ostracod assemblages in Osaka Bay, one of the most polluted marine areas in the world.

Benthic foraminiferal and ostracod assemblages from surface sediment in Osaka Bay, collected in 2004, were analyzed to characterize the distribution of the modern foraminiferal and ostracod assemblages. The results suggested that

low diversity community, typically characterized by exclusive dominance of three eutrophication-tolerant foraminiferal species (*Ammonia beccarii*, *Eggerella advena*, and *Trochammina hadai*), is found in the inner part of the bay, where eutrophication is serious.

Four sediment cores, which obtained along a transect from mouth of the Yodo River (a source of nutrient input) to middle part of Osaka Bay, were examined to trace the development process of modern low diversity community. The records of foraminiferal and ostracod assemblages in the cores clearly documented the development of low diversity community after the urbanization in the early part of the 20th century. Many foraminiferal and ostracod species that are unable to tolerate low-oxygen conditions decreased and three eutrophication-tolerant foraminiferal species increased after the early part of the 20th century, when Japan's industrial revolution started. As a consequence, low diversity community developed rapidly in the inner part of the bay since that time. Low diversity community most progressed as a result of severe hypoxia and food increase by eutrophication during the high economic growth period (~1970's). A sewage treatment program for the Seto Inland Sea was enacted in 1970's in order to reduce the organic pollutant loads. Coincident with the commencing period of the program, the density of eutrophication-tolerant foraminiferal species decreased, but no major changes in the foraminiferal and ostracod assemblages occurred. This study provides new information regarding the use of foraminifera and ostracod as paleoecological indicators of anthropogenic environmental/ecosystem changes.

Annual variations of Oxygen Deficient Water Mass (ODW) in Jinhae Bay

Chang Yong MOON^{*}, In-Cheol LEE & Yoo Jihn OH

Department of Ocean Engineering, Pukyong National University, Busan 608-737, Korea

* E-mail: liong83@nate.com

As a basic study to improve oxygen deficient water mass (ODW), this study analyzed the spatial and temporal variation of the water quality and investigated the characteristics of ODW in Jinhae bay, from 1989 to 2006. By observing the annual variation of water qualities, the severe polluted area was found, and the effects of a special management sea area in Masan bay and the construction of Pusan Newport on the water